



The Thoracic Society
of Australia & New Zealand
LEADERS IN LUNG HEALTH

Thoracic Society of Australia and New Zealand

PHYSIOTHERAPY FOR CYSTIC FIBROSIS IN AUSTRALIA AND NEW ZEALAND

Clinical Practice Guideline

April 2016

Button, B. M., Wilson, C., Dentice, R., Cox, N. S., Middleton, A., Tannenbaum, E., Bishop, J., Cobb, R., Burton, K., Wood, M., Moran, F., Black, R., Bowen, S., Day, R., Depiazzi, J., Doiron, K., Doumit, M., Dwyer, T., Elliot, A., Fuller, L., Hall, K., Hutchins, M., Kerr, M., Lee, A. L., Mans, C., O'Connor, L., Steward, R., Potter, A., Rasekaba, T., Scoones, R., Tarrant, B., Ward, N., West, S., White, D., Wilson, L., Wood, J., and Holland, A. E. (2016) Physiotherapy for cystic fibrosis in Australia and New Zealand: A clinical practice guideline. *Respirology*, 21: 656–667. doi: [10.1111/resp.12764](https://doi.org/10.1111/resp.12764).

The extended version of the clinical practice guidelines presented in this paper can be found in the online supplementary information. Any reference to the online supplementary information should cite this **Respirology** paper as the primary publication.

<http://onlinelibrary.wiley.com/doi/10.1111/resp.12764/full>

FOREWORD

This Clinical Practice Guideline has been written by physiotherapists who are experienced in the management of cystic fibrosis in Australia and New Zealand.

Chairperson: Associate Professor Brenda Button
Senior Clinician Physiotherapist
Alfred Hospital and Monash
University, Melbourne

Editor: Professor Anne Holland
Clinical Chair, Physiotherapy
La Trobe University / Alfred Health,
Melbourne

Brenda Button Alfred Hospital and Monash University, Melbourne

Christine Wilson Royal Children's Hospital, Queensland

Ruth Dentice Royal Prince Alfred Hospital, NSW

Narelle Cox La Trobe University, Victoria

Anna Middleton Children's Hospital at Westmead, NSW

Esta-Lee Tannenbaum Royal Children's Hospital, Victoria

Jenny Bishop Westmead Hospital, NSW

Robyn Cobb The Prince Charles Hospital, Queensland

Kate Burton The Prince Charles Hospital, Queensland

Michelle Wood The Prince Charles Hospital, Queensland

Fiona Moran The Royal Children's Hospital, Victoria

Ryan Black Lady Cilento Children's Hospital, Brisbane

Summar Bowen Princess Margaret Hospital, Western Australia

Rosie Day John Hunter Children's Hospital, NSW

Julie Depiazzi Princess Margaret Hospital, Western Australia

Katherine Doiron Bond University/ Pindara Private Hospital, Qld

Michael Doumit Sydney Children's Hospital, NSW

Tiffany Dwyer Central Clinical School, Sydney Medical School, University of Sydney; Royal Prince Alfred Hospital, NSW.

Alison Elliot The Children's Hospital at Westmead, NSW

Louise Fuller Alfred Hospital, Victoria

Kathleen Hall The Prince Charles Hospital, Queensland

Matthew Hutchins Mater Health Services

Melinda Kerr Royal Children's Hospital, Victoria

Annemarie Lee Alfred Hospital, Victoria

Christina Mans Waikato DHB, Hamilton NZ

Lauren O'Connor Gold Coast University Hospital, Queensland

Ranjana Steward Alfred Hospital, Victoria

Angela Potter Women's & Children's Hospital, SA

Tshepo Rasekaba Alfred Hospital, Victoria

Rebecca Scoones Starship Children's Health, New Zealand

Ben Tarrant Alfred Hospital, Victoria

Nathan Ward Royal Adelaide Hospital, SA

Samantha West Westmead Hospital, NSW

Dianne White Royal Adelaide Hospital, South Australia

Lisa Wilson Alfred Hospital, Victoria

Jamie Wood Sir Charles Gairdner Hospital and Institute for Respiratory Health, Western Australia

Anne Holland La Trobe University / Alfred Health, Melbourne

ABSTRACT

Physiotherapy management is a key element of care for people with cystic fibrosis (CF) throughout the lifespan. Although considerable evidence exists to support physiotherapy management of CF, there is documented variation in practice. The aim of this guideline is to optimise the physiotherapy management of people with CF in Australia and New Zealand. A systematic review of the literature in key areas of physiotherapy practice for CF was undertaken. Recommendations were formulated based on National Health and Medical Research Council (Australia) guidelines and considered the quality, quantity and level of the evidence; the consistency of the body of evidence; the likely clinical impact; and applicability to physiotherapy practice in Australia and New Zealand. A total of 30 recommendations were made for airway clearance therapy; inhalation therapy; exercise assessment and training; musculoskeletal management; management of urinary incontinence; managing the newly diagnosed patient with CF; delivery of non-invasive ventilation; and physiotherapy management before and after lung transplantation. These recommendations can be used to underpin the provision of evidence-based physiotherapy care to people with CF in Australia and New Zealand.

Key words: Cystic fibrosis; physiotherapy, airway clearance; inhalation therapy; exercise

Short title: Physiotherapy Guidelines for Cystic Fibrosis



BACKGROUND

The aim of this Clinical Practice Guideline is to optimise physiotherapy management of people with CF in Australia and New Zealand. Recommendations for key areas of physiotherapy management are provided, including airway clearance therapy; inhalation therapy; exercise and musculoskeletal management. This Clinical Practice Guideline builds on a previous Consensus Statement for physiotherapy management of CF in Australia¹ and is informed by an evaluation of its uptake and impact.²

METHODS

All physiotherapists who were regularly caring for people with CF across paediatric and adult settings in Australia and New Zealand were invited to participate in guideline development. A systematic literature search was undertaken for each topic area up until June 2014, using MEDLINE, CINAHL, EMBASE and PEDro. Data from each included study were extracted into an evidence table by one reviewer and checked by a second reviewer. Quality was graded according to the NHMRC evidence hierarchy.³ Recommendations were formulated based on the quality, quantity and level of the evidence; the consistency of the body of evidence; the likely clinical impact; and generalisability and applicability to physiotherapy practice in Australia and New Zealand.³ Areas of importance to physiotherapy practice, but with insufficient evidence to make recommendations, were highlighted. Updates of each section were prepared by a writing group and circulated to all authors for comment and revision. The draft document was offered to stakeholders for comment, including CF physicians, CF consumers, allied health professionals and physiotherapists who were not part of the writing group.

This Guideline provides recommendations for clinical physiotherapy practice and a summary of the evidence that underpins them. More details regarding the evidence underpinning the recommendations and application of the physiotherapy techniques can be found in the Online Supplement.

AIRWAY CLEARANCE TECHNIQUES

The aim of airway clearance techniques (ACTs) is to clear sputum from the airway, in order to optimise respiratory status and slow disease progression. Airway clearance techniques are frequently described as a 'cornerstone' of CF treatment.⁴ A Cochrane review concluded that ACTs have short-term beneficial effects on mucus transport in CF.⁵ One uncontrolled study evaluated the effects of withdrawing airway clearance for three weeks and found a detrimental effect on pulmonary function.⁶ Due to ethical concerns regarding withholding such a well-established treatment, it is considered unlikely that more robust controlled trials of the long term impact of ACTs in CF will be conducted.⁷

A number of effective ACTs are available. The active cycle of breathing technique (ACBT) consists of breathing control, thoracic expansion exercises and forced expirations.⁸ The ACBT is effective for clearance of respiratory secretions⁹ with results comparable to other widely used ACTs.¹⁰ The ACBT can be performed by all patients who can follow instructions and is useful in all stages of disease. Positive expiratory pressure (PEP) therapy is defined as breathing against a positive expiratory pressure of 10 – 20 cmH₂O¹¹ using a mask or mouthpiece. A Cochrane review concluded that PEP was equally effective as other forms of ACT and that patients may prefer PEP.¹² Oscillating PEP combines oscillation of airflow with PEP, in order to loosen secretions.¹³ In a one-year RCT comparing oscillating PEP with PEP in children with CF, greater deterioration in pulmonary function and more hospitalisations were seen in the oscillating PEP group.¹⁴ However, a more recent 1-year study in adults showed no difference in lung function between groups randomly assigned to PEP or oscillating PEP.¹⁵ Autogenic drainage (AD) uses controlled breathing to achieve the highest possible airflow in different generations of bronchi.¹⁶ Short-term studies have shown that AD is as effective as postural drainage and percussion,¹⁷ oscillating PEP¹⁸ and ACBT.¹⁹ In a long term comparative study in adolescents with CF, AD was as effective as postural drainage, and

participants showed strong preference for AD.²⁰ Several systematic reviews note that no single ACT is superior,^{10, 12, 21} such that treatment choices should be individualised.

Postural drainage involves use of gravity to drain mucus from the lungs. A number of studies have demonstrated provocation of gastro-oesophageal reflux (GOR) during head-down tilted postural drainage in infants, children and adolescents with CF.²²⁻²⁴ Two additional studies did not reproduce these results in infants, with no differences in GOR between modified and traditional postural drainage,^{25, 26} however the head-down position utilised was not as steep, older infants were studied and they avoided the prone head down tilted position.²⁵ Other potential adverse effects of postural drainage with head-down tilt include increased dyspnoea²⁷ and oxyhaemoglobin desaturation.²⁸

Modified postural drainage involves positioning without use of head-down tilt.²² In a five year follow up of infants randomised to either standard or modified postural drainage, the modified group had fewer radiological changes and significantly better lung function at six years of age.²⁹ In a short-term adult study comparing treatment in head down versus horizontal positions, there was no difference in amount of sputum expectorated, but patients reported fewer side effects in horizontal positions.²⁷

Physical exercise that increases minute ventilation leads to the mobilisation of pulmonary secretions and enhances airway clearance.³⁰⁻³³ Some people with mild CF lung disease use exercise together with forced expiration and coughing as a stand-alone ACT. Others with more extensive lung disease and larger volumes of sputum use exercise as an adjunct to a formal ACT regimen. Physiological effects of exercise include reduced mechanical impedance of sputum, enhanced expiratory flow rates and inducement of coughing.³⁴⁻³⁸ A meta-analysis including three trials found that the addition of exercise to ACTs significantly increased FEV₁ compared to ACT alone.³⁹ Whether exercise can be used as an alternative to formal ACTs is less clear, with conflicting results across trials.^{31, 32, 38, 40}

- 1. Airway clearance techniques should be performed across the lifespan in CF (C)**
- 2. The ACBT is an effective form of airway clearance and can be used by people with acute and chronic lung disease independently or in conjunction with other ACTs (B)**
- 3. PEP therapy, oscillating PEP and autogenic drainage are effective forms of airway clearance which can be performed independently (B)**
- 4. Postural drainage in head-down positions should not be used routinely in infants with CF (B) or in patients of any age with known or suspected GOR (C). Modified postural drainage is recommended in infants and young children where active participation in airway clearance therapy is not possible (B)**
- 5. Physical exercise may be used to reduce mechanical impedance of sputum (B), achieve short term improvements in pulmonary function (A) and improve ease of expectoration (B)**

INHALATION THERAPY

Inhalation therapy is an important treatment for CF respiratory disease. Effective inhalation therapy is integral to the success of ACTs and vice versa; as a result, physiotherapists should be adequately skilled in delivery of inhalation therapy in order to maximise the effectiveness of both treatments.

The main determinants of deposition pattern for nebulised medications are the breathing pattern during inhalation, droplet size and age/condition of the lung.⁴¹ A slower breath results in a more desirable peripheral deposition pattern, improved homogeneity of deposition and increased overall drug deposition.⁴² Slow steady breaths with occasional deep breaths have traditionally been recommended to promote improved deposition.⁴³ However, the specific device being utilised may determine the optimal breathing pattern.⁴⁴ Given the varying physico-chemical behaviours of the nebulised medications in CF, it is important to use a nebuliser/compressor combination which is effective for the specific preparation.⁴¹

Nebulised medication should be taken via a mouthpiece to maximise delivery of the drug to the airways and avoid nasal filtration.⁴⁵ Exceptions are in young children who may be unable to use a mouthpiece effectively, or



where sinuses are a target of therapy, or those with acute dyspnoea. Bronchodilators should be delivered by metered dose inhaler and spacer⁴¹ except where patients are too dyspnoeic or are unable to follow instructions. Metered dose inhalers with spacer should be used for administration of inhaled corticosteroids. Patients should be encouraged to rinse their mouth with water and gargle afterwards to reduce the risk of thrush.⁴⁵

Combining ACTs and inhalation therapy can reduce the time-related burden of care in CF. However studies have reported conflicting results of this practice on lung deposition, bronchodilation, dyspnoea, cough and mucus production.⁴⁶⁻⁴⁸ The combination of PEP with inhalation therapy (hypertonic or isotonic saline, salbutamol) is commonly used by physiotherapists. Some also combine inhalation therapy with positioning and breathing techniques such as ACBT or AD. There is insufficient research investigating the combination of inhalation therapy and ACTs to make recommendations regarding this practice.

The optimal timing of inhalation of dornase alfa in relation to ACTs has been investigated in a Cochrane systematic review.⁴⁹ Meta-analysis showed that inhalation of dornase alfa after ACT had similar effects on FEV₁, FVC and quality of life compared to inhalation prior to ACT. However, FEF₂₅ was significantly better with dornase alfa inhalation before ACT, based on the pooled data from two small studies in children with well preserved pulmonary function.^{50, 51} A longer time interval between administration of dornase alfa and ACT is more effective than inhalation immediately preceding ACT.⁵² There appears to be no detrimental effects on sleep quality or nocturnal cough if this time interval is extended so that dornase alfa is administered before bedtime.⁵³ In the absence of strong evidence to indicate that one regimen is better than another, the timing of dornase alfa inhalation in relation to ACT or time of day can be based on pragmatic reasons or individual preference.

The timing of hypertonic saline inhalation in relation to ACT (before, during or after) does not appear to have a substantial effect on lung function after a single treatment session.⁵⁴ However, participants were more satisfied with the entire treatment session when hypertonic saline was inhaled before or during ACT and perceived these timing regimens as more effective.

- 6. Where possible, nebulised medication should be taken via a mouthpiece (C)**
- 7. To optimise dose delivery and treatment time, inhalation technique should be adapted to the device being used, including consideration of body position and concurrent ACTs (C)**
- 8. Bronchodilators should be delivered by metered dose inhaler unless there is clinical need for nebulisation (C)**
- 9. Metered dose inhalers with spacer should be used for the administration of inhaled corticosteroids. (B)**
- 10. Hypertonic saline may be administered before or during airway clearance techniques (B)**

EXERCISE ASSESSMENT AND TRAINING

Measures of exercise capacity predict survival in children and adults with CF^{55, 56} and those with better physical fitness have better quality of life.⁵⁷ Structured exercise programs for people with CF improve fitness, thoracic mobility, maintain bone mineral density and may slow the rate of pulmonary decline.⁵⁸⁻⁶⁴ A study involving over 200 people with CF conducted over nine years showed that patients with higher levels of physical activity in daily life (encompassing all activities, not just exercise programs) had a slower rate of FEV₁ decline than those who were less active.⁶⁵

Exercise testing enables evaluation of exercise capacity, functional capacity, response to treatment and disease progression.^{55, 66-68} For physiotherapists, exercise tests also provide the basis for exercise prescription. Commonly



used field exercise tests in CF are the 6-minute walk test, the modified shuttle walk test and the 3-minute step test. More details regarding the conduct and choice of exercise test are provided in the full document online (<http://www.thoracic.org.au/professional-information/position-papers-guidelines/cystic-fibrosis/>). The six-minute walk test is considered a particularly important measure during preparation for lung transplantation in CF, providing guidance regarding the timing of referral.⁶⁹

Both aerobic training and anaerobic training are beneficial in CF.⁵⁸ Aerobic training results in improved maximum exercise capacity, strength and quality of life.^{70, 71} Anaerobic training has positive effects on lactate levels, peak power⁷² and fat-free mass.⁷¹ Both types of exercise may have positive effects on pulmonary function.^{71, 73} Exercise programs have beneficial effects both during admission for acute exacerbations^{38, 71, 74} and for the stable outpatient.^{72, 73, 75} It is unclear whether home-based, unsupervised training programs are equally as effective as supervised programs.

The ideal exercise prescription for people with CF has not been established. In the absence of specific guidelines, aerobic exercise prescription should follow the same principles as those used in healthy individuals⁷⁶ and patients with other chronic respiratory diseases.⁷⁷ Exercise training should occur on at least three days (preferably five or more days) per week; have a duration of 30 minutes per session; and increase heart rate to 75% of maximum heart rate. A combination of aerobic and resistance training is required to achieve maximum benefits.⁵⁸

Patients with CF may exhibit exercise-induced oxygen desaturation during training, even when pulmonary function is well preserved.⁷⁸ Supplemental oxygen during training increases exercise duration;⁷⁹ whether this improves clinical outcomes is not clear. Supplemental oxygen is frequently used during training in patients whose oxygen saturation falls below 90% during exercise.⁸⁰

- 11. Exercise is recommended for people with CF throughout the lifespan (B)**
- 12. An exercise test should be considered to assess response to therapy in the inpatient and outpatient settings and as an assessment tool in the prescription of exercise training programs (C)**
- 13. A six minute walk test should be performed as part of the initial assessment for lung transplantation (C).**
- 14. Exercise prescription should be tailored to the individual and comply with recommended exercise guidelines (B)**
- 15. Supplemental oxygen should be considered during training in patients with severe exercise-induced desaturation (C)**

MUSCULOSKELETAL COMPLICATIONS OF CF

Musculoskeletal manifestations of CF arise as a result of multi-factorial abnormalities in bone mineralization, altered respiratory mechanics and muscular imbalance secondary to pulmonary disease. Between 43% and 94% of individuals with CF experience spinal pain⁸¹⁻⁹², which occurs across the spectrum of disease severity. Musculoskeletal pain in CF is associated with decreased quality of life, increased respiratory symptoms, sleep disturbance, anxiety, depression, and a reduced ability to perform ACTs and exercise.^{82, 86, 88-95}

People with CF have multiple risk factors for inadequate bone mineralisation.⁹⁶⁻¹⁰⁰ Longitudinal studies have shown that bone gains during puberty are decreased in CF adolescents compared to healthy controls, resulting in decreased attainment of peak bone mass.^{97, 98, 101-106} A meta-analysis reported that in CF the prevalence of osteoporosis and osteopenia were 23.5% and 38% respectively.¹⁰⁷ Individuals with more severe lung disease, decreased physical activity and low aerobic capacity had the lowest bone mineral density, higher prevalence of vertebral fractures and more severe kyphosis.¹⁰⁸



Weight bearing exercise is the most effective non-pharmacological method to improve bone mineral density in the healthy population by stimulating bone accretion, preventing bone loss and improving bone structural qualities.¹⁰⁶ As the foundation of bone health begins in childhood, and there is some evidence of sustained benefit from early vigorous physical activity,^{109, 110} children and adolescents should engage in high impact weight bearing exercise for 30 minutes three times a week. The pre-pubertal and early pubertal years are particularly important to help maximise peak bone mass^{101, 111, 112} as approximately one quarter of peak bone mass is gained in the two years around the pubertal growth spurt.^{111, 113}

Changes in muscle strength, length and neuromuscular recruitment have been demonstrated in CF. Reduced lean muscle mass is associated with malabsorption and deconditioning.^{114, 115} Peripheral muscle impairment is also noted in response to systemic inflammation and lack of moderate to vigorous physical activity.^{116, 117} In CF, resistance training programs of moderate to high intensity and variable duration (19 days – 12 months) have produced significant leg strength gains.¹¹⁸⁻¹²⁰ Strength training for children is still a novel area with protocols not clearly defined; this is an area for future research.¹²⁰

Physiotherapists require the skills to manage a diverse range of musculoskeletal complications including CF related arthropathy, sports injuries, spinal pain, and pain associated with coughing. Early reports regarding the use of manual therapy and exercise in the management of pain and restriction are encouraging.^{121, 122} Patients with CF-related arthropathy require rheumatological management, which may include physiotherapy interventions targeting pain reduction and muscle strengthening.

- 1. A musculoskeletal assessment should be included at annual review from approximately age eight (pre puberty). Earlier assessment is warranted if pain or functional impairment is reported, or bone mineral density risk highlighted (C)**
- 2. Regular physical activity, including weightbearing exercise, should be encouraged throughout the lifespan in order to optimise bone density (C)**
- 3. Strength training programs should be prescribed in order to optimise muscle mass (B)**

PHYSIOTHERAPY MANAGEMENT OF THE COMPLEX PATIENT

Cystic fibrosis is a complex multi-system disease and patients often experience complications and comorbidities that have implications for physiotherapy management. These include haemoptysis, pneumothorax, CF-related diabetes and pregnancy.

There are no published data regarding physiotherapy management of patients with haemoptysis or pneumothorax, however guidelines based on expert opinion are available.¹²³ When haemoptysis is present, the physiotherapist aims to maintain adequate airway clearance and exercise regimens whilst promoting vessel healing and minimising the risk of re-bleeding. When a pneumothorax is present, physiotherapists aim to ensure that adequate airway clearance continues, whilst minimising the amount of positive pressure generated inside the patient's lungs. Both of these situations may require alteration to usual ACTs.

CF-related diabetes (CFRD) is a frequent comorbidity in CF, occurring in 5-30% of patients. The American Diabetes Association Clinical Care Guidelines for CFRD¹²⁴ state that people with CFRD should perform moderate aerobic exercise for at least 150 minutes per week; should monitor blood glucose levels (BGLs) before vigorous physical activity; and may need to consume extra carbohydrate or alter their insulin dose (level of evidence –expert opinion). During periods of acute illness or courses of corticosteroids, blood sugar levels and insulin requirements may be altered and more careful monitoring required.¹²⁴

Many pregnancy-related physiological changes have implications for optimal physiotherapy care, although little research is available. Women with CF are encouraged to approach pregnancy with a regular ACT routine. Head-down tilted postural drainage should be avoided, along with any ACTs that exacerbate nausea. Upright sitting is



usually the most comfortable position for airway clearance. Modifications to exercise programs may be required to accommodate musculoskeletal, respiratory and cardiac changes. Maintenance of adequate hydration during exercise should be emphasised. It is important to consider the need for domestic support during pregnancy and afterwards, to provide sufficient time for regular airway clearance, inhalation therapy and exercise.¹²⁵

PHYSIOTHERAPY MANAGEMENT OF CONTINENCE

The reported prevalence of urinary incontinence in girls and women with CF ranges from 22% to 74%¹²⁶⁻¹³² in comparison with 13% in healthy women of similar age.¹³³ There is limited literature in adult males with CF, with a reported prevalence of 8-15%, compared to 7.5% in healthy men.^{132, 134, 135} It is not known whether the cause of urinary incontinence in CF is chronic cough; loading of the pelvic floor during ACTs, coughing and physical exercise; or underlying structural differences. People with CF and incontinence report increased anxiety, depression and a negative impact on quality of life.^{135, 136}

Screening for incontinence should be part of routine physiotherapy care for both male and female patients. Treatment of urinary incontinence in women with CF by a continence physiotherapist with exercise, electrical stimulation, biofeedback and bladder training results in improvements in pelvic floor strength, reduction in leakage and improvement in quality of life.¹³⁷ Positive outcomes have also been demonstrated with surgical correction of severe urinary incontinence in women with CF.¹³⁸ In order to optimise pelvic floor function, patients should be taught to perform ACTs in positions that maintain a neutral lumbar spine,¹³⁹ with addition of perineal support in those with urinary incontinence.

- 19. Women with CF and symptoms of stress urinary incontinence should be taught rehabilitative strength and endurance exercises to provide better control of the pelvic floor (C)**
- 20. Men and women with CF should be screened for symptoms of stress urinary incontinence (C)**
- 21. Airway clearance should take place in postures that maintain a neutral lumbar spine, to optimize pelvic floor function (C)**

PHYSIOTHERAPY MANAGEMENT OF THE NEWLY DIAGNOSED PATIENT

Newly diagnosed infants and their families should meet with the CF multi-disciplinary team soon after diagnosis.¹⁴⁰ ¹⁴¹ The role of physiotherapy in ACTs, exercise and active play should be explained, demonstrated and practiced.¹⁴² ¹⁴³ Treatment of infants should follow the usual guidelines for physiotherapy in CF, consisting of five modified postural drainage positions performed 1-2 times daily as appropriate.^{22, 24, 29} In each position percussion or thoracic compressions should be performed for 3-5 minutes. Other techniques such as infant PEP or assisted autogenic drainage can also be introduced.¹⁴⁴ Normal developmental play and prone lying should be encouraged as the first steps towards an active physical lifestyle and routine.^{144, 145}

Adults and children with a new diagnosis of CF usually have milder disease than those diagnosed in infancy.¹⁴² Some newly diagnosed adults are very well; therefore physical exercise, huffing and coughing may be appropriate



as stand-alone ACTs.³⁴ Those patients with established lung disease should be taught appropriate ACTs and exercise regimens, and educated about the role of nebulised drugs in their treatment.

- 22. Treatment for newly diagnosed infants may include percussion for 3-5 minutes in each of 5 modified postural drainage positions (B) and daily age-appropriate physical play (C)**
- 23. Physiotherapy treatment for the newly diagnosed child and adult should include regular physical exercise (B); other forms of airway clearance therapy should be added as required (C)**

NON-INVASIVE VENTILATION FOR CF

A number of descriptive studies report the successful use of NIV to stabilise patients with CF and acute respiratory failure,¹⁴⁶⁻¹⁵² with reduced hypercapnia, respiratory rate and dyspnoea. Although NIV does not reverse the respiratory deterioration inherent in end-stage disease, it may allow the patient to be stabilised for long enough for donor lungs to become available for transplantation. The use of NIV for CF patients who are not awaiting lung transplantation has also been reported¹⁵¹ where it may be useful for palliation of dyspnoea in end-stage disease.

Hypoxia and hypercapnia occur commonly during sleep in moderate to severe CF^{153, 154} and may result in daytime respiratory failure.¹⁵⁵ Positive short-term effects of NIV during sleep in CF have been reported.^{156, 157} Longer-term outcomes of NIV for chronic respiratory failure may include improvements in daytime PaCO₂, reduction in the number of days spent in hospital and improvement in symptoms.^{158, 159} In a randomised controlled trial, domiciliary NIV over a 6 week period resulted in significant improvements in quality of life, respiratory symptom scores, dyspnoea, nocturnal ventilation and increased exercise performance.¹⁶⁰

Airway clearance techniques are onerous for patients who are unwell, due to increased ventilatory demand,¹⁶¹ alterations in gas exchange^{28, 162} and dyspnoea.²⁷ Two randomised crossover trials report that a single session of NIV can unload the respiratory muscles during ACTs in both adults and children with CF, resulting in decreased dyspnoea and less desaturation.^{162, 163} Similar effects have been observed with the application of NIV during exercise, including improved ventilation, reduced desaturation and increased functional walking performance.¹⁶⁴ This may be useful in patients bridging to transplantation, in whom maintenance of exercise capacity is an important goal.

Non-invasive ventilation delivers air at high flow rates and low relative humidity, which may overwhelm the capacity of the upper airway mucosa to warm and humidify inspired air.¹⁶⁵ Humidity levels during NIV are low enough to cause airway drying.¹⁶⁶ This is of greatest concern in patients with excessive secretions, who are at high risk of sputum retention. Hence, consideration should be given to heated humidification when NIV is used in CF.

- 24. Non-invasive ventilation should be considered in all patients with acute respiratory failure who are listed for transplantation (C)**
- 25. In patients with symptomatic nocturnal ventilatory failure, a trial of nocturnal NIV may be undertaken (B)**
- 26. NIV is a useful adjunct to airway clearance in patients with severe disease in whom dyspnoea and fatigue limits effective airway clearance (B)**
- 27. NIV may be a useful adjunct to exercise in patients with severe disease where dyspnoea and fatigue contribute to deconditioning and limit effective training (B)**
- 28. Heated humidification should be incorporated into the circuit for all applications of NIV in CF (C)**

PHYSIOTHERAPY AND LUNG TRANSPLANTATION

There are many systemic features of CF which have the potential to impact on lung transplant suitability and outcomes, including skeletal muscle weakness¹⁶⁷ and poor bone health.⁶³ Most adult transplant centres offer dedicated pre-operative exercise training classes for transplant candidates, in order to optimise physical fitness and strength. A recent large, retrospective study which included 70 people with CF¹⁶⁸ showed that 6-minute walk distance was well maintained from listing to transplantation in those who undertook thrice-weekly supervised exercise training. Furthermore, those with a greater 6-minute walk distance prior to transplantation had a shorter hospital stay post-transplantation.

Exercise rehabilitation is an established therapy for lung transplant recipients.¹⁶⁹ Although studies in CF are uncontrolled, three months of post transplant rehabilitation has been associated with improvements in functional exercise capacity, strength and quality of life in adults¹⁷⁰ and children.¹⁷¹ A recent randomised controlled trial of three months of rehabilitation in lung transplantation recipients with other respiratory disorders, performed immediately following hospital discharge, showed significant improvements in daily physical activity, quadriceps force and exercise performance at one year following transplantation.¹⁷² The content of post transplantation rehabilitation programs generally includes aerobic and resistance exercise, performed at least three times per week.¹⁶⁹ Although most rehabilitation takes place in a group setting, patients with resistant organisms may require isolation from other immune suppressed patients.

29. Patients with CF should undertake an exercise program designed to optimise their physical function while on the transplant waiting list (C)

30. Patients with CF who have undergone lung transplantation should participate in a formal, supervised rehabilitation program post-operatively (B)

END OF LIFE CARE

There is no published literature specifically addressing physiotherapy treatment in the terminal stages of CF. However many people with CF die of respiratory failure¹⁷³ and physiotherapists are often involved with provision of end of life care. The aims of physiotherapy treatment will be influenced by whether the patient is actively waiting for transplantation. Care should focus on comfort and dignity and be tailored to each patient's goals and values.¹⁷⁴ Minimising the work of breathing during ACTs is an important consideration in those approaching the end of life. Some patients may require therapist-assisted ACTs such as percussion or thoracic compressions. Airway clearance may be continued even in the palliative stage if the patient finds it beneficial to relieve symptoms. Comfort measures such as soft tissue massage and positioning can also be considered.

INFECTION CONTROL

Respiratory pathogens have a significant impact on morbidity and mortality in CF¹⁷⁵ and good infection control practices are critical to preventing their transmission. Segregation and cohorting of inpatients and outpatients according to respiratory organisms are now routinely practiced.^{176, 177} Physiotherapists are encouraged to be familiar with their local infection control policies. Due to considerable variation in local policies, formal recommendations are not provided here.

Both *Pseudomonas aeruginosa* and *B. cepacia* may be spread in droplet form by coughing and can survive on dry surfaces for a number of days.¹⁷⁸⁻¹⁸⁰ There is also potential for airborne transmission.¹⁸¹ *B. cepacia* has been isolated from the hospital rooms and hands of patients following airway clearance¹⁸²⁻¹⁸⁴ and the outside surfaces



of sputum cups.¹⁸² These findings reinforce the need to segregate patients whilst performing ACTs and inhalation therapy, as well as the importance of hand washing. Stethoscopes should be cleaned with alcohol wipes between patients.¹⁸⁵

Bacterial contamination of home nebulisers has been documented and sharing equipment has been associated with transmission of *B. cepacia*.¹⁸⁶⁻¹⁸⁸ Under no circumstances should any respiratory equipment be shared between patients with CF. There is no consensus regarding the use of gloves, gowns and masks during physiotherapy treatment in CF. Physiotherapists should consult their local infection control policy with regard to when these measures are required.

Coughing is common during exercise and droplet spread of organisms is possible. These droplets may be transmitted within one metre of an infected patient.¹⁷⁸ It has been reported that contamination can still occur between 1 and 2 metres, albeit with lower probability (1.7%).¹⁸⁹ As a result, patients with different organisms, or in different cohorts, do not exercise together. When people considered suitable for cohorting are sharing the gym, universal precautions should be practised. Patients should be educated to maintain a two metre distance from other patients at all times; hands should be washed on entering and leaving the gym; and patients should be taught to wipe down all exercise equipment with an alcohol-based solution before and after use.

PHYSIOTHERAPY SERVICES FOR CF

There is no published research regarding the optimum structure of physiotherapy services for people with CF. For inpatients, expert clinical opinion suggests that physiotherapy assessment and treatment starts on the day of hospital admission.¹⁹⁰ The physiotherapy treatment plan should address inhalation therapy, ACT and exercise. Patients admitted with an acute exacerbation with increased and/or retained secretions will need to carry out more frequent ACT sessions than their baseline daily regimen. The number of treatments will range from two to three or more treatments in 24 hours. A graduated physical exercise program incorporating cardiorespiratory exercise should be commenced as soon as possible.

Patients attending the outpatient department of a Cystic Fibrosis Service should have access to a physiotherapist with expertise in CF management at each clinic visit. It is suggested that each patient be assessed 3-6 monthly so that their physiotherapy program can be reviewed and optimized. Complex patients may require more frequent and detailed review. A formal annual review by the CF team, including physiotherapy review, has been advocated for people with CF.^{190, 191}

ACKNOWLEDGEMENTS

The contribution of the authors was as follows - Chair of working group: BMB; Editor: AEH; Writing group leaders: CW, RD, NC, AM, ET, BJ, RC, KM, MW, FM, RB; Writing group members: SB, RD, RDay, JD, KD, MD, TD, AE, LF, KH, MH, NK, MK, AL, CM, LO, RS, AP, TR, RS, BT, NW, SW, DW, LW, JW, SW. The authors would like to thank Ms Nardia Roberston, Dr Audrey Tierney, Dr Dominic Keating and Dr Kathy Stiller for their helpful comments and review of this document.

DISCLOSURE STATEMENT

Development of the 2015 Clinical Practice Guideline was partially funded by a grant from Cystic Fibrosis Australia to support the editorial role of Dr Holland and was used to fund a research assistant. Development of the 2008 Consensus Statement (of which this guideline is an update and extension) was funded in part by unrestricted educational grants from Roche Pharmaceuticals and Solvay Pharmaceuticals. This funding was used to support travel of the writing group to the consensus conference in Melbourne in September 2006 and to assist dissemination by hard copy and CD ROM. A grant from Cystic Fibrosis Australia supported the editorial role of Dr Holland. The funders had no role in the decision to develop the document, nor in its preparation or writing. Declarations of interest were made by each author according to the policies of the Thoracic Society of Australia and New Zealand. BB received conference support from Roche and Pharmaxis and is on the Allied Health Advisory Committees of Pharmaxis, Novartis and Vertex. RB (2013) and RC received conference support from Novartis. NC and RDay (2007;2009) received conference support from Roche. AM received conference support from Solvay, Technipro and Pharmaxis. FM received a Technipro Scholarship in 2012. KM received conference support from Pharmaxis (2012) and Novartis (2013). NW received conference support Roche (2007) and Pharmaxis (2012). CW received meeting support from Pharmaxis. JW received conference support from Roche (2009), Pharmaxis (2012) and Novartis (2013); an Abbott Products scholarship (2011); and Bronchitol workshop (2012). MW received advisory board consultancy fees from Vertex (2014) and travel support to attend clinical trials investigator meetings from Vertex (2010 – 2014), PTC (2014), Aradigm (2014).



REFERENCES

1. Physiotherapy for Cystic Fibrosis in Australia: A Consensus Statement. <http://www.thoracic.org.au/documents/papers/physiotherapyforcf.pdf>. Accessed 20th March 2015
2. Holland AE, Button BM. Physiotherapy for cystic fibrosis in Australia: Knowledge and acceptance of the Consensus Statement recommendations. *Respirology*. 2013; **18**: 652-6.
3. National Health and Medical Research Council. 2009. NHMRC additional levels of evidence and grades for recommendations for developers of guidelines. https://www.nhmrc.gov.au/_files_nhmrc/file/guidelines/developers/nhmrc_levels_grades_evidence_120423.pdf. Accessed: 20th March 2015.
4. Yankaskas JR, Marshall BC, Sufian B, Simon RH, Rodman D. Cystic Fibrosis Adult Care: Consensus Conference Report. *Chest*. 2004; **125**: 1S-39.
5. van der Schans C, Prasad A, Main E. Chest physiotherapy compared to no chest physiotherapy for cystic fibrosis. *Cochrane Database of Systematic Reviews*. 2003; **3**: 3.
6. Desmond KJ, Schwenk WF, Thomas E, Beaudry PH, Coates AL. Immediate and long-term effects of chest physiotherapy in patients with cystic fibrosis. *J Paediatr*. 1983; **103**: 538-42.
7. Prasad A. Physiotherapy in cystic fibrosis. *Journal of the Royal Society of Medicine*. 2000; **93**: 27-36.
8. Pryor J. Active cycle of breathing techniques. In: Mcllwaine M, Van Ginderdeuren F, (eds.) *Physiotherapy in the treatment of cystic fibrosis*. 3rd ed. International Physiotherapy Group/ Cystic Fibrosis, 2009; 5-7.
9. Pryor J, Webber B, Hodson M, Batter J. Evaluation of the forced expiration technique as an adjunct to postural drainage in the treatment of cystic fibrosis. *Br Med J*. 1979; **2**: 417-8.
10. McKoy NA, Saldanha IJ, Odelola OA, Robinson KA. Active cycle of breathing technique for cystic fibrosis. *Cochrane Database Syst Rev*. 2012; **12**: CD007862.
11. Lannefors L, Eriksson L. Positive Expiratory Pressure (PEP). In: Mcllwaine M, Van Ginderdeuren F, (eds.) *Physiotherapy in the treatment of cystic fibrosis*. 3rd ed. International Physiotherapy Group/ Cystic Fibrosis, 2009; 12-4.
12. Elkins MR, Jones A, van der Schans C. Positive expiratory pressure physiotherapy for airway clearance in people with cystic fibrosis. *Cochrane Database Syst Rev*. 2006: CD003147.
13. Althaus P. Oscillating PEP - Flutter therapy. In: Mcllwaine M, Van Ginderdeuren F, (eds.) *Physiotherapy in the treatment of cystic fibrosis*. 3rd ed. International Physiotherapy Group/ Cystic Fibrosis, 2009; 18-22.
14. Mcllwaine PM, Wong LT, Peacock D, Davidson AG. Long-term comparative trial of positive expiratory pressure versus oscillating positive expiratory pressure (flutter) physiotherapy in the treatment of cystic fibrosis. *J Pediatr*. 2001; **138**: 845-50.
15. Newbold M, Tullis E, Corey M, Ross B, Brooks D. The flutter device versus the PEP mask in the treatment of adults with cystic fibrosis. *Physiotherapy Canada*. 2005; **57**: 199-207.
16. Chevallier J. Autogenic Drainage (AD). In: Mcllwaine M, Van Ginderdeuren F, (eds.) *Physiotherapy in the treatment of cystic fibrosis (CF)*. International Physiotherapy Group/ Cystic Fibrosis, 2009; 8-9.
17. Giles DR, Wagener JS, Accurso FJ, Butler-Simon N. Short-term effects of postural drainage with clapping vs autogenic drainage on oxygen saturation and sputum recovery in patients with cystic fibrosis. *Chest*. 1995; **108**: 952-4.
18. App EM, Kieselmann R, Reinhardt D, Lindemann H, Dasgupta B, King M, Brand P. Sputum rheology changes in cystic fibrosis lung disease following two different types of physiotherapy: flutter vs autogenic drainage. *Chest*. 1998; **114**: 171-7.
19. Miller S, Hall DO, Clayton CB, Nelson R. Chest physiotherapy in cystic fibrosis: a comparative study of autogenic drainage and the active cycle of breathing techniques with postural drainage. *Thorax*. 1995; **50**: 165-9.
20. Mcllwaine M, Wong LT, Chilvers M, Davidson GF. Long-term comparative trial of two different physiotherapy techniques; postural drainage with percussion and autogenic drainage, in the treatment of cystic fibrosis. *Pediatr Pulmonol*. 2010; **45**: 1064-9.
21. Morrison L, Agnew J. Oscillating devices for airway clearance in people with cystic fibrosis. *Cochrane Database Syst Rev*. 2009: CD006842.
22. Button BM, Heine RG, Catto-Smith AG, Phelan PD, Olinsky A. Postural drainage and gastro-oesophageal reflux in infants with cystic fibrosis. *Arch Dis Child*. 1997; **76**: 148-50.
23. Vandenplas Y, Dierickx A, Blecker U, Lanciers S, Deneyer M. Esophageal pH monitoring data during chest physiotherapy. *J Pediatr Gastroenterol Nutr*. 1991; **13**: 23-6.
24. Button BM, Heine RG, Catto-Smith AG, Phelan PD. Postural drainage in cystic fibrosis: is there a link with gastro-oesophageal reflux? *J Paediatr Child Health*. 1998; **34**: 330-4.
25. Phillips GE, Pike SE, Rosenthal M, Bush A. Holding the baby: head downwards positioning for physiotherapy does not cause gastro-oesophageal reflux. *Eur Respir J*. 1998; **12**: 954-7.
26. Doumit M, Krishnan U, Jaffe A, Belessis Y. Acid and non-acid reflux during physiotherapy in young children with cystic fibrosis. *Pediatr Pulmonol*. 2012; **47**: 119-24.
27. Cecins N, Jenkins S, Pengelly J, Ryan G. The active cycle of breathing techniques - to tip or not to tip? *Respir Med*. 1999; **93**: 660-5.
28. McDonnell T, McNicholas WT, FitzGerald MX. Hypoxaemia during chest physiotherapy in patients with cystic fibrosis. *Ir J Med Sci*. 1986; **155**: 345-8.



29. Button BM, Heine RG, Catto-Smith AG, Olinsky A, Phelan PD, Ditchfield MR, Story I. Chest physiotherapy in infants with cystic fibrosis: to tip or not? A five-year study. *Pediatr Pulmonol.* 2003; **35**: 208-13.
30. Hebestreit A, Kersting U, Basler B, Jeschke R, Hebestreit H. Exercise inhibits epithelial sodium channels in patients with cystic fibrosis. *Am J Respir Crit Care Med.* 2001; **164**: 443-6.
31. Lannefors L, Wollmer P. Mucus clearance with three chest physiotherapy regimes in cystic fibrosis: a comparison between postural drainage, PEP and physical exercise. *Eur Respir J.* 1992; **5**: 748-53.
32. Salh W, Bilton D, Dodd M, Webb AK. Effect of exercise and physiotherapy in aiding sputum expectoration in adults with cystic fibrosis. *Thorax.* 1989; **44**: 1006-8.
33. Zach M, Oberwaldner B, Hausler F. Cystic fibrosis: physical exercise versus chest physiotherapy. *Arch Dis Child.* 1982; **57**: 587-9.
34. Dwyer TJ, Alison JA, McKeough ZJ, Daviskas E, Bye PT. Effects of exercise on respiratory flow and sputum properties in patients with cystic fibrosis. *Chest.* 2011; **139**: 870-7.
35. Bilton D, Dodd ME, Abbot JV, Webb AK. The benefits of exercise combined with physiotherapy in the treatment of adults with cystic fibrosis. *Respir Med.* 1992; **86**: 507-11.
36. Loughlin GM, Cota KA, Taussig LM. The relationship between flow transients and bronchial lability in cystic fibrosis. *Chest.* 1981; **79**: 206-10.
37. Macfarlane PI, Heaf D. Changes in airflow obstruction and oxygen saturation in response to exercise and bronchodilators in cystic fibrosis. *Pediatr Pulmonol.* 1990; **8**: 4-11.
38. Cerny FJ. Relative effects of bronchial drainage and exercise for in-hospital care of patients with cystic fibrosis. *Phys Ther.* 1989; **69**: 633-9.
39. Thomas J, Cook, D.J., Brooks, D. Chest physical therapy management of patients with cystic fibrosis: a meta-analysis. *Am J Respir Crit Care Med.* 1995; **151**: 846-50.
40. Reix P, Aubert F, Werck-Gallois MC, Toutain A, Mazzocchi C, Moreux N, Bellon G, Rabilloud M, Kassai B. Exercise with incorporated expiratory manoeuvres was as effective as breathing techniques for airway clearance in children with cystic fibrosis: a randomised crossover trial. *J Physiother.* 2012; **58**: 241-7.
41. Boe J, Dennis JH, O'Driscoll BR, Bauer TT, Carone M, Dautzenberg B, Diot P, Heslop K, Lannefors L. European Respiratory Society Guidelines on the use of nebulizers. *Eur Respir J.* 2001; **18**: 228-42.
42. Laube BL, Jashnani R, Dalby RN, Zeitlin PL. Targeting aerosol deposition in patients with cystic fibrosis: effects of alterations in particle size and inspiratory flow rate. *Chest.* 2000; **118**: 1069-76.
43. Nebulizer therapy. Guidelines. British Thoracic Society Nebulizer Project Group. *Thorax.* 1997; **52 Suppl 2**: S4-24.
44. Heijerman H, Westerman E, Conway S, Touw D, Doring G. Inhaled medications and inhalation devices for lung disease in patients with cystic fibrosis: a European consensus. *Journal of Cystic Fibrosis* 2009; **8**: 295-315.
45. Laube BL, Janssens HM, de Jongh FH, Devadason SG, Dhand R, Diot P, Everard ML, Horvath I, Navalesi P, Voshaar T, Chrystyn H, European Respiratory S, International Society for Aerosols in M. What the pulmonary specialist should know about the new inhalation therapies. *Eur Respir J.* 2011; **37**: 1308-31.
46. Laube BL, Geller DE, Lin TC, Dalby RN, Diener-West M, Zeitlin PL. Positive expiratory pressure changes aerosol distribution in patients with cystic fibrosis. *Respir Care.* 2005; **50**: 1438-44.
47. Frischknecht-Christensen E, Norregaard O, Dahl R. Treatment of bronchial asthma with terbutaline inhaled by conespace combined with positive expiratory pressure mask. *Chest.* 1991; **100**: 317-21.
48. Stites SW, Perry GV, Peddicord T, Cox G, McMillan C, Becker B. Effect of high-frequency chest wall oscillation on the central and peripheral distribution of aerosolized diethylene triamine penta-acetic acid as compared to standard chest physiotherapy in cystic fibrosis. *Chest.* 2006; **129**: 712-7.
49. Dentice R, Elkins M. Timing of dornase alfa inhalation for cystic fibrosis. *Cochrane Database of Systematic Reviews.* 2011: CD007923.
50. Anderson P, Morton J. Evaluation of two different timings of Pulmozyme nebulisation in relation to chest physiotherapy in children with cystic fibrosis [abstract]. *Journal of Cystic Fibrosis.* 2009; **8**: S74.
51. van der Giessen LJ, de Jongste JC, Gosselink R, Hop WC, Tiddens HA. RhDNase before airway clearance therapy improves airway patency in children with CF. *Pediatr Pulmonol.* 2007; **42**: 624-30.
52. Wilson CJ, Robbins LJ, Murphy JM, Chang AB. Is a longer time interval between recombinant human deoxyribonuclease (dornase alfa) and chest physiotherapy better?: A multi-center, randomized crossover trial. *Pediatr Pulmonol.* 2007; **42**: 1110-6.
53. van der Giessen LJ, Gosselink R, Hop WC, Tiddens HA. Recombinant human DNase nebulisation in children with cystic fibrosis: before bedtime or after waking up? *Eur Respir J.* 2007; **30**: 763-8.
54. Dentice RL, Elkins MR, Bye PT. Adults with cystic fibrosis prefer hypertonic saline before or during airway clearance techniques: a randomised crossover trial. *J Physiother.* 2012; **58**: 33-40.
55. Nixon PA, Orenstein DM, Kelsey SF, Doershuk CF. The prognostic value of exercise testing in patients with cystic fibrosis. *N Engl J Med.* 1992; **327**: 1785-8.
56. Pianosi P, Leblanc J, Almudevar A. Peak oxygen uptake and mortality in children with cystic fibrosis. *Thorax.* 2005; **60**: 50-4.
57. Orenstein DM, Franklin BA, Doershuk CF, Hellerstein HK, Germann KJ, Horowitz JG, Stern RC. Exercise conditioning and cardiopulmonary fitness in cystic fibrosis. The effects of a three-month supervised running program. *Chest.* 1981; **80**: 392-8.
58. Bradley J, Moran F. Physical training for Cystic fibrosis. *Cochrane Database of Systematic Reviews.* 2008; **2008**.
59. Hebestreit H, Kieser S, Junge S, Ballmann M, Hebestreit A, Schindler C, Schenk T, Posselt HG, Kriemler S. Long-term effects of a partially supervised conditioning programme in cystic fibrosis. *European Respiratory Journal.* 2010; **35**: 578-83.



60. Elbasan B, Tunali N, Duzgun I, Ozcelik U. Effects of chest physiotherapy and aerobic exercise training on physical fitness in young children with cystic fibrosis. *Ital J Pediatr.* 2012; **38**: 2.
61. Hulzebos H, Snieder H, van der Et J, Helders PJ, Takken T. High-intensity interval training in an adolescent with cystic fibrosis: a physiological perspective. *Physiother Theory Pract.* 2011; **27**: 231-7.
62. Shoemaker M, Hurt H, Arndt L. The evidence regarding exercise training in the management of cystic fibrosis: a systematic review. *Cardiopulm Phys Ther J.* 2008; **19**: 75-83.
63. Tejero Garcia S, Giraldez Sanchez M, Cejudo P, Quintana Gallego E, Dapena J, Carcia Jimenez R, Can Luis P, Gomez de Terreros I. Bone health, daily physical activity, and exercise tolerance in patients with cystic fibrosis. *Chest.* 2011; **140**: 475-81..
64. Paranjape S, Barnes L, Carson K, von Berg K, Hoosen H, Mogayzel P. Exercise improves lung function and habitual activity in children with cystic fibrosis. *J Cyst Fibros.* 2012; **11**: 18-23. Epub 2011 Sep 3.
65. Schneiderman JE, Wilkes DL, Atenafu EG, Nguyen T, Wells GD, Alarie N, Tullis E, Lands LC, Coates AL, Corey M, Ratjen F. Longitudinal relationship between physical activity and lung health in patients with cystic fibrosis. *Eur Respir J.* 2014; **43**: 817-23.
66. Bradley J, Howard J, Wallace E, Elborn S. Reliability, repeatability, and sensitivity of the modified shuttle test in adult cystic fibrosis. *Chest.* 2000; **117**: 1666-71.
67. Webb AK, Dodd ME, Moorcroft J. Exercise and cystic fibrosis. *J R Soc Med.* 1995; **88 Suppl 25**: 30-6.
68. Alison JA, Donnelly PM, Lennon M, Parker S, Torzillo P, Mellis C, Bye PT. The effect of a comprehensive, intensive inpatient treatment program on lung function and exercise capacity in patients with cystic fibrosis. *Phys Ther.* 1994; **74**: 583-91; discussion 91-3.
69. Kadikar A, Maurer J, Kesten S. The six-minute walk test: a guide to assessment for lung transplantation. *J Heart Lung Transplant.* 1997; **16**: 313-9.
70. Schmidt AM, Jacobsen U, Bregnballe V, Olesen HV, Ingemann-Hansen T, Thastum M, Oluf Schiøtz P. Exercise and quality of life in patients with cystic fibrosis: A 12-week intervention study. *Physiother Theory Pract.* 2011; **27**: 548-56. Epub 2011 Jul 3.
71. Selvadurai HC, Blimkie CJ, Meyers N, Mellis CM, Cooper PJ, Van Asperen PP. Randomized controlled study of in-hospital exercise training programs in children with cystic fibrosis. *Pediatr Pulmonol.* 2002; **33**: 194-200.
72. Klijn PH, Oudshoorn A, van der Ent CK, van der Net J, Kimpen JL, Helders PJ. Effects of anaerobic training in children with cystic fibrosis: a randomized controlled study. *Chest.* 2004; **125**: 1299-305.
73. Schneiderman-Walker J, Pollock SL, Corey M, Wilkes DD, Canny GJ, Pedder L, Reisman JJ. A randomized controlled trial of a 3-year home exercise program in cystic fibrosis. *J Pediatr.* 2000; **136**: 304-10.
74. Kuys S, Hall K, Peasey M, Wood M, Cobb R, Bell S. Gaming console exercise and cycle or treadmill exercise provide similar cardiovascular demand in adults with cystic fibrosis: a randomised crossover trial. *J Physiother.* 2011; **57**: 35-40.
75. Moorcroft AJ, Dodd ME, Morris J, Webb AK. Individualised unsupervised exercise training in adults with cystic fibrosis: a 1 year randomised controlled trial. *Thorax.* 2004; **59**: 1074-80.
76. American College of Sports Medicine. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness and flexibility in healthy adults. *Med Sci Sports Exerc.* 1998; **30**: 975-91.
77. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, Hill K, Holland AE, Lareau SC, Man WD, Pitta F, Sewell L, Raskin J, Bourbeau J, Crouch R, Franssen FM, Casaburi R, Vercoulen JH, Vogiatzis I, Gosselink R, Clini EM, Effing TW, Maltais F, van der Palen J, Troosters T, Janssen DJ, Collins E, Garcia-Aymerich J, Brooks D, Fahy BF, Puhon MA, Hoogendoorn M, Garrod R, Schols AM, Carlin B, Benzo R, Meek P, Morgan M, Rutten-van Molken MP, Ries AL, Make B, Goldstein RS, Dowson CA, Brozek JL, Donner CF, Wouters EF. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med.* 2013; **188**: e13-64.
78. Bradley J, Howard J, Wallace E, Elborn S. Validity of a modified shuttle test in adult cystic fibrosis. *Thorax.* 1999; **54**: 437-9.
79. Mallory GB, Fullmer JJ, Vaughan DJ. Oxygen therapy for cystic fibrosis. *Cochrane Database Syst Rev.* 2005: CD003884.
80. Pulmonary Rehabilitation. British Thoracic Society Standards of Care Subcommittee on Pulmonary Rehabilitation. *Thorax.* 2001; **56**: 827-34.
81. Ravilly S, Robinson W, Suresh S, Wohl ME, Berde CB. Chronic pain in cystic fibrosis. *Pediatrics.* 1996; **98**: 741-7.
82. Festini F, Ballarin S, Codamo T, Doro R, Loganés C. Prevalence of pain in adults with cystic fibrosis. *J Cyst Fibros.* 2004; **3**: 51-7.
83. Botton E, Saraux A, Laselve H, Jousse S, Le Goff P. Musculoskeletal manifestations in cystic fibrosis. *Joint Bone Spine.* 2003; **70**: 327-35.
84. Parasa RB, Maffulli N. Musculoskeletal involvement in cystic fibrosis. *Bulletin (Hospital for Joint Diseases) New York, NY.* 1999; **58**: 37-44.
85. Tattersall R, Walshaw MJ. Back pain in adult patients with CF: effect of chronic steroid use and physiotherapy. *Journal of Cystic Fibrosis.* 2001; **1**.
86. Flume PA, Ciolino J, Gray S, Lester MK. Patient-reported pain and impaired sleep quality in adult patients with cystic fibrosis. *J Cyst Fibros.* 2009; **8**: 321-5.
87. Hayes M, Yaster M, Haythornthwaite JA, Riekert KA, McMillan KN, White E, Mogayzel PJ, Jr., Lechtzin N. Pain is a common problem affecting clinical outcomes in adults with cystic fibrosis. *Chest.* **140**: 1598-603.
88. Kelemen L, Lee AL, Button BM, Presnell S, Wilson JW, Holland AE. Pain Impacts on Quality of Life and Interferes with Treatment in Adults with Cystic Fibrosis. *Physiother Res Int.* 2012; **17**: 132-41.
89. Ravilly S, Robinson W, Suresh S, Wohl ME, Berde CB. Chronic pain in cystic fibrosis. *Pediatrics.* 1996; **98**: 741-7.
90. Sawicki GS, Sellers DE, Robinson WM. Self-reported physical and psychological symptom burden in adults with cystic fibrosis. *J Pain Symptom Manage.* 2008; **35**: 372-80.



91. Sermet-Gaudelus I, De Villartay P, de Dreuzy P, Clairicia M, Vrielynck S, Canoui P, Kirzsenbaum M, Singh-Mali I, Agrario L, Salort M, Charron B, Dusser D, Lenoir G, Hubert D. Pain in children and adults with cystic fibrosis: a comparative study. *J Pain Symptom Manage*. 2009; **38**: 281-90.
92. Stenekes SJ, Hughes A, Gregoire MC, Frager G, Robinson WM, McGrath PJ. Frequency and self-management of pain, dyspnea, and cough in cystic fibrosis. *J Pain Symptom Manage*. 2009; **38**: 837-48.
93. Koh JL, Harrison D, Palermo TM, Turner H, McGraw T. Assessment of acute and chronic pain symptoms in children with cystic fibrosis. *Pediatr Pulmonol*. 2005; **40**: 330-5.
94. Eksterowicz N. Pain management in cystic fibrosis. *Paediatric Pulmonology*. 2000; **Supplement 20**: 114-5.
95. Chastain DC, Cook AJ. Chronic pain in CF: associated beliefs and behaviours. *Pediatric Pulmonology*. 2000; **Supplement 20**: 116-7.
96. Grey V, Atkinson S, Drury D, Casey L, Ferland G, Gundberg C, Lands LC. Prevalence of low bone mass and deficiencies of vitamins D and K in pediatric patients with cystic fibrosis from 3 Canadian centers. *Pediatrics*. 2008; **122**: 1014-20.
97. Javier RM, Jacquot J. Bone disease in cystic fibrosis: what's new? *Joint, bone, spine : revue du rhumatisme*. 2011; **78**: 445-50.
98. Haworth CS. Impact of cystic fibrosis on bone health. *Current opinion in pulmonary medicine*. 2010; **16**: 616-22.
99. Sparks AA, McGee SJ, Boone CE, Neuringer IP, Jones SK, Aris RM. 'Old' bones in young bodies: the tale of cystic fibrosis. *Curr Opin Endocrinol Diabetes Obes*. 2009; **16**: 407-14.
100. Sermet-Gaudelus I, Castanet M, Retsch-Bogart G, Aris RM. Update on cystic fibrosis-related bone disease: a special focus on children. *Paediatric Respiratory Reviews*. 2009; **10**: 134-42.
101. Sermet-Gaudelus I, Bianchi ML, Garabedian M, Aris RM, Morton A, Hardin DS, Elkin SL, Compston JE, Conway SP, Castanet M, Wolfe S, Haworth CS. European cystic fibrosis bone mineralisation guidelines. *Journal of Cystic Fibrosis*. 2011; **10 Suppl 2**: S16-23.
102. Buntain HM, Schluter PJ, Bell SC, Greer RM, Wong JC, Batch J, Lewindon P, Wainwright CE. Controlled longitudinal study of bone mass accrual in children and adolescents with cystic fibrosis. *Thorax*. 2006; **61**: 146-54.
103. Buntain HM, Greer RM, Schluter PJ, Wong JC, Batch JA, Potter JM, Lewindon PJ, Powell E, Wainwright CE, Bell SC. Bone mineral density in Australian children, adolescents and adults with cystic fibrosis: a controlled cross sectional study. *Thorax*. 2004; **59**: 149-55.
104. Bianchi ML, Romano G, Saraifoger S, Costantini D, Limonta C, Colombo C. BMD and body composition in children and young patients affected by cystic fibrosis. *J Bone Miner Res*. 2006; **21**: 388-96.
105. Curran DR, McArdle JR, Talwalkar JS. Diabetes mellitus and bone disease in cystic fibrosis. *Semin Respir Crit Care Med*. 2009; **30**: 514-30.
106. Hind K, Truscott JG, Conway SP. Exercise during childhood and adolescence: a prophylaxis against cystic fibrosis-related low bone mineral density? Exercise for bone health in children with cystic fibrosis. *Journal of Cystic Fibrosis*. 2008; **7**: 270-6.
107. Paccou J, Zeboulon N, Combescure C, Gossec L, Cortet B. The prevalence of osteoporosis, osteopenia, and fractures among adults with cystic fibrosis: a systematic literature review with meta-analysis. *Calcif Tissue Int*. 2010; **86**: 1-7.
108. Tejero Garcia S, Giraldez Sanchez MA, Cejudo P, Quintana Gallego E, Dapena J, Garcia Jimenez R, Cano Luis P, Gomez de Terreros I. Bone health, daily physical activity, and exercise tolerance in patients with cystic fibrosis. *Chest*. 2011; **140**: 475-81.
109. Janz KF, Letuchy EM, Eichenberger Gilmore JM, Burns TL, Torner JC, Willing MC, Levy SM. Early physical activity provides sustained bone health benefits later in childhood. *Medicine and Science in Sports and Exercise*. 2010; **42**: 1072-8.
110. Baxter-Jones AD, Kontulainen SA, Faulkner RA, Bailey DA. A longitudinal study of the relationship of physical activity to bone mineral accrual from adolescence to young adulthood. *Bone*. 2008; **43**: 1101-7.
111. Barry DW, Kohrt WM. Exercise and the preservation of bone health. *J Cardiopulm Rehabil Prev*. 2008; **28**: 153-62.
112. Body JJ, Bergmann P, Boonen S, Boutsen Y, Bruyere O, Devogelaer JP, Goemaere S, Hollevoet N, Kaufman JM, Milisen K, Rozenberg S, Reginster JY. Non-pharmacological management of osteoporosis: a consensus of the Belgian Bone Club. *Osteoporosis International*. 2011; **22**: 2769-88.
113. Boreham CA, McKay HA. Physical activity in childhood and bone health. *Br J Sports Med*. 2011; **45**: 877-9.
114. Lands L, Desmond KJ, Demizio D, Pavilanis A, Coates AL. The effects of nutritional status and hyperinflation on respiratory muscle strength in children and young adults. *Am Rev Respir Dis*. 1990; **141**: 1506-9.
115. Lands LC, Heigenhauser GJ, Jones NL. Respiratory and peripheral muscle function in cystic fibrosis. *Am Rev Respir Dis*. 1993; **147**: 865-9.
116. Lamhonwah AM, Bear CE, Huan LJ, Kim Chiaw P, Ackerley CA, Tein I. Cystic fibrosis transmembrane conductance regulator in human muscle: Dysfunction causes abnormal metabolic recovery in exercise. *Ann Neurol*. 2010; **67**: 802-8.
117. Burtin C, Van Remoortel H, Vrijnsen B, Langer D, Colpaert K, Gosselink R, Decramer M, Dupont L, Troosters T. Impact of exacerbations of cystic fibrosis on muscle strength. *Respir Res*. 2013; **14**: 46.
118. Orenstein DM, Hovell MF, Mulvihill M, Keating KK, Hofstetter CR, Kelsey S, Morris K, Nixon PA. Strength vs aerobic training in children with cystic fibrosis: a randomized controlled trial. *Chest*. 2004; **126**: 1204-14.
119. Selvadurai HC, Blimkie CJ, Meyers N, Mellis CM, Cooper PJ, Van Asperen PP. Randomized controlled study of in-hospital exercise training programs in children with cystic fibrosis. *Pediatric Pulmonology*. 2002; **33**: 194-200.
120. van Doorn N. Exercise programs for children with cystic fibrosis: a systematic review of randomized controlled trials. *Disability and Rehabilitation*. 2010; **32**: 41-9.
121. Lee A, Holdsworth M, Holland A, Button B. The immediate effect of musculoskeletal physiotherapy techniques and massage on pain and ease of breathing in adults with cystic fibrosis. *Journal of Cystic Fibrosis*. 2009; **8**: 79-81.



122. Sandsund CA, Roughton M, Hodson ME, Pryor JA. Musculoskeletal techniques for clinically stable adults with cystic fibrosis: a preliminary randomised controlled trial. *Physiotherapy*. 2011; **97**: 209-17.
123. Flume PA, Mogayzel Jr PJ, Robinson KA, Rosenblatt RL, Quittell L, Marshall BC, Committee CPGFPT. Cystic Fibrosis Pulmonary Guidelines: Pulmonary Complications: Hemoptysis and Pneumothorax. *Am J Respir Crit Care Med*. 2010; **182**: 298-306.
124. Moran A, Brunzell C, Cohen RC, Katz M, Marshall BC, Onady G, Robinson KA, Sabadosa KA, Stecenko A, Slovis B. Clinical care guidelines for cystic fibrosis-related diabetes: a position statement of the American Diabetes Association and a clinical practice guideline of the Cystic Fibrosis Foundation, endorsed by the Pediatric Endocrine Society. *Diabetes Care*. 2010; **33**: 2697-708.
125. Johannesson M. Effects of pregnancy on health: certain aspects of importance for women with cystic fibrosis. *J Cyst Fibros*. 2002; **1**: 9-12.
126. Cornacchia M, Zenorini A, Perobelli S, Zanolta L, Mastella G, Braggion C. Prevalence of urinary incontinence in women with cystic fibrosis. *BJU international*. 2001; **88**: 44-8.
127. White D, Stiller K, Roney F. The prevalence and severity of symptoms of incontinence in adult cystic fibrosis patients. *Physiotherapy Theory and Practice*. 2000; **16**: 35-42.
128. Orr A, McVean RJ, Webb AK, Dodd ME. Questionnaire survey of urinary incontinence in women with cystic fibrosis. *BMJ*. 2001; **322**: 1521.
129. Nixon GM, Glazner JA, Martin JM, Sawyer SM. Urinary incontinence in female adolescents with cystic fibrosis. *Pediatrics*. 2002; **110**: e22.
130. Moran F, Bradley JM, Boyle L, Elborn JS. Incontinence in adult females with cystic fibrosis: a Northern Ireland survey. *International Journal of Clinical Practice*. 2003; **57**: 182-3.
131. Vella M, Cartwright R, Cardozo L, Parsons M, Madge S, Burns Y. Prevalence of incontinence and incontinence-specific quality of life impairment in women with cystic fibrosis. *Neurourology and Urodynamics*. 2009; **28**: 986-9.
132. Browne WJ, Wood CJ, Desai M, Weller PH. Urinary incontinence in 9-16 year olds with cystic fibrosis compared to other respiratory conditions and a normal group. *Journal of Cystic Fibrosis*. 2009; **8**: 50-7.
133. Chiarelli P, Brown W, McElduff P. Leaking urine: prevalence and associated factors in Australian women. *Neurourology and Urodynamics*. 1999; **18**: 567-77.
134. Burge AT, Holland AE, Sherburn M, Wilson JW, Cox N, Rasekaba TM, McAleer R, Morton J, Button BM. Prevalence and impact of incontinence in adult men with cystic fibrosis. *Respirology*. 2011; **16**: 57.
135. Gumery L, Hodgson G, Humphries N, Sheldon J, Stableforth D, Mackenzie W, Honeybourne D, Hawkins G. The prevalence of urinary incontinence in the adult male population of a regional cystic fibrosis centre. *Journal of Cystic Fibrosis*. 2002; **1**: S173.
136. Nankivell G, Caldwell P, Follett J. Urinary incontinence in adolescent females with cystic fibrosis. *Paediatric Respiratory Reviews*. 2010; **11**: 95-9.
137. Button BM, Sherburn M, Chase J, Stillman B, Wilson J. Effect of a Three Months Physiotherapeutic Intervention on Incontinence in Women with Chronic Cough Related to Cystic Fibrosis and COPD. *Pediatric Pulmonology*. 2005; **Suppl 28**: A369.
138. Helm JM, Langman H, Dodd ME, Ahluwalia A, Jones AM, Webb AK. A novel solution for severe urinary incontinence in women with cystic fibrosis. *Journal of Cystic Fibrosis*. 2008; **7**: 501-4.
139. Sapsford RR, Richardson CA, Stanton WR. Sitting posture affects pelvic floor muscle activity in parous women: an observational study. *Aust J Physiother*. 2006; **52**: 219-22.
140. Sawyer SM, Glazner JA. What follows newborn screening? An evaluation of a residential education program for parents of infants with newly diagnosed cystic fibrosis. *Pediatrics*. 2004; **114**: 411-6.
141. Merelle ME, Huisman J, Alderden-van der Vecht A, Taat F, Bezemer D, Griffioen RW, Brinkhorst G, Dankert-Roelse JE. Early versus late diagnosis: psychological impact on parents of children with cystic fibrosis. *Pediatrics*. 2003; **111**: 346-50.
142. Kerem E, Conway S, Elborn S, Heijerman H. Standards of care for patients with cystic fibrosis: a European consensus. *J Cyst Fibros*. 2005; **4**: 7-26.
143. Physiotherapy in the treatment of cystic fibrosis. 2009. https://www.ecfs.eu/ipg_cf/booklet. Accessed: January 15th 2015.
144. Lannefors L, Button BM, McLwaine M. Physiotherapy in infants and young children with cystic fibrosis: current practice and future developments. *J R Soc Med*. 2004; **97 Suppl 44**: 8-25.
145. Dennersten U, Lannefors L, Högglund P, Hellberg K, Johansson H, Lagerkvist AL, Ortfelt M, Sahlberg M, Eriksson L. Lung function in the aging Swedish cystic fibrosis population. *Respir Med*. 2009; **103**: 1076-82.
146. Hodson ME, Madden BP, Steven MH, Tsang VT, Yacoub MH. Non-invasive mechanical ventilation for cystic fibrosis patients--a potential bridge to transplantation. *Eur Respir J*. 1991; **4**: 524-7.
147. Piper AJ, Parker S, Torzillo PJ, Sullivan CE, Bye PT. Nocturnal nasal IPPV stabilizes patients with cystic fibrosis and hypercapnic respiratory failure. *Chest*. 1992; **102**: 846-50.
148. Padman R, Lawless S, Von Nessen S. Use of BiPAP by nasal mask in the treatment of respiratory insufficiency in pediatric patients: preliminary investigation. *Pediatr Pulmonol*. 1994; **17**: 119-23.
149. Caronia CG, Silver P, Nimkoff L, Gorvov J, Quinn C, Sagy M. Use of bilevel positive airway pressure (BIPAP) in end-stage patients with cystic fibrosis awaiting lung transplantation. *Clinical Pediatrics*. 1998; **37**: 555-9.
150. Sprague K, Graff G, Tobias DJ. Noninvasive ventilation in respiratory failure due to cystic fibrosis. *South Med J*. 2000; **93**: 954-61.
151. Madden BP, Kariyawasam H, Siddiqi AJ, Machin A, Pryor JA, Hodson ME. Noninvasive ventilation in cystic fibrosis patients with acute or chronic respiratory failure. *Eur Respir J*. 2002; **19**: 310-3.
152. Flight WG, Shaw J, Johnson S, Webb AK, Jones AM, Bentley AM, Bright-Thomas RJ. Long-term non-invasive ventilation in cystic fibrosis - experience over two decades. *Journal of Cystic Fibrosis*. 2012; **11**: 187-92.
153. Muller NL, Francis PW, Gurwitz D, Levison H, Bryan AC. Mechanism of hemoglobin desaturation during rapid-eye-movement sleep in normal subjects and in patients with cystic fibrosis. *Am Rev Respir Dis*. 1980; **121**: 463-9.



154. Tepper RS, Skatrud JB, Dempsey JA. Ventilation and oxygenation changes during sleep in cystic fibrosis. *Chest*. 1983; **84**: 388-93.
155. Olson EJ, Simon PM. Sleep-wake cycles and the management of respiratory failure. *Curr Opin Pulm Med*. 1996; **2**: 500-6.
156. Gozal D. Nocturnal ventilatory support in patients with cystic fibrosis: comparison with supplemental oxygen. *Eur Respir J*. 1997; **10**: 1999-2003.
157. Milross MA, Piper AJ, Norman M, Becker HF, Willson GN, Grunstein RR, Sullivan CE, Bye PT. Low-flow Oxygen and Bilevel Ventilatory Support. Effects on ventilation during sleep in cystic fibrosis. *Am J Respir Crit Care Med*. 2001; **163**: 129-34.
158. Hill AT, Edenborough FP, Cayton RM, Stableforth DE. Long-term nasal intermittent positive pressure ventilation in patients with cystic fibrosis and hypercapnic respiratory failure (1991-1996). *Respir Med*. 1998; **92**: 523-6.
159. Granton JT, Shapiro C, Kesten S. Noninvasive nocturnal ventilatory support in advanced lung disease from cystic fibrosis. *Respir Care*. 2002; **47**: 675-81.
160. Young AC, Wilson JW, Kotsimbos TC, Naughton MT. Randomised placebo controlled trial of non-invasive ventilation for hypercapnia in cystic fibrosis. *Thorax*. 2008; **63**: 72-7.
161. Williams MT, Parsons DW, Frick RA, Ellis ER, Martin AJ, Giles SE, Grant ER. Acute respiratory infection in patients with cystic fibrosis with mild pulmonary impairment: comparison of two physiotherapy regimens. *Aust J Physiother*. 2001; **47**: 227-36.
162. Holland AE, Denehy L, Ntoumenopoulos G, Naughton MT, Wilson JW. Non-invasive ventilation assists chest physiotherapy in adults with acute exacerbations of cystic fibrosis. *Thorax*. 2003; **58**: 880-4.
163. Fauroux B, Boule M, Lofaso F, Zerah F, Clement A, Harf A, Isabey D. Chest physiotherapy in cystic fibrosis: improved tolerance with nasal pressure support ventilation. *Pediatrics*. 1999; **103**: 1-9.
164. Lima CA, de Andrade Ade F, Campos SL, Brandao DC, Fregonezi G, Mourato IP, Aliverti A, de Britto MC. Effects of noninvasive ventilation on treadmill 6-min walk distance and regional chest wall volumes in cystic fibrosis: Randomized controlled trial. *Respir Med*. 2014; **108**: 1460-8.
165. Holland AE, Denehy L, Buchan CA, Wilson JW. Efficacy of a heated passover humidifier during noninvasive ventilation: a bench study. *Respir Care*. 2007; **52**: 38-44.
166. Wiest GH, Foerst MD, Fuchs FS, Schmelzer AH, Hahn EG, Ficker JH. In vivo efficacy of two heated humidifiers used during CPAP-therapy for obstructive sleep apnea under various environmental conditions. *Sleep*. 2001; **24**: 435-40.
167. Troosters T, Langer D, Vrijzen B, Segers J, Wouters K, Janssens W, Gosselink R, Decramer M, Dupont L. Skeletal muscle weakness, exercise tolerance and physical activity in adults with cystic fibrosis. *European Respiratory Journal*. 2009; **33**: 99-106.
168. Li M, Mathur S, Chowdhury NA, Helm D, Singer LG. Pulmonary rehabilitation in lung transplant candidates. *J Heart Lung Transplant*. 2013; **32**: 626-32.
169. Wickerson L, Mathur S, Brooks D. Exercise training after lung transplantation: a systematic review. *The Journal of Heart and Lung Transplantation*. 2010; **29**: 497-503.
170. Munro PE, Holland AE, Bailey M, Button BM, Snell GI. Pulmonary rehabilitation following lung transplantation. *Transplant Proc*. 2009; **41**: 292-5.
171. Deliva RD, Hassall A, Manthot C, Solomon M, McCrindle BW, Dipchand AI. Effects of an acute, outpatient physiotherapy exercise program following pediatric heart or lung transplantation. *Pediatric Transplantation*. 2012; **16**: 879-86.
172. Langer D, Burtin C, Schepers L, Ivanova A, Verleden G, Decramer M, Troosters T, Gosselink R. Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am J Transplant*. 2012; **12**: 1584-92.
173. Davis PB. Clinical pathophysiology and manifestations of lung disease. In: Yankaskas JR, (ed.) *Cystic Fibrosis in Adults*. 1st edition ed. Lippincott-Raven, Philadelphia, 1999; 45-67.
174. Tonelli MR. End-of-life care in cystic fibrosis. *Curr Opin Pulm Med*. 1998; **4**: 332-6.
175. Ledson MJ, Gallagher MJ, Jackson M, Hart CA, Walshaw MJ. Outcome of Burkholderia cepacia colonisation in an adult cystic fibrosis centre. *Thorax*. 2002; **57**: 142-5.
176. 2012. Infection control guidelines for cystic fibrosis patients and carers. <http://www.cysticfibrosis.org.au/cfa/infection-control>. Accessed: 15th January 2015 2015.
177. Saiman L, Siegel JD, LiPuma JJ, Brown RF, Bryson EA, Chambers MJ, Downer VS, Fliege J, Hazle LA, Jain M, Marshall BC, O'Malley C, Pattee SR, Potter-Bynoe G, Reid S, Robinson KA, Sabadosa KA, Schmidt HJ, Tullis E, Webber J, Weber DJ. Infection Prevention and Control Guideline for Cystic Fibrosis: 2013 Update. *Infection Control and Hospital Epidemiology*. 2014; **35**.
178. Saiman L, Siegel J. Infection control in cystic fibrosis. *Clinical Microbiology Reviews*. 2004; **17**: 57-71.
179. Zimakoff J, Hoiby N, Rosendal K, Guilbert JP. Epidemiology of Pseudomonas aeruginosa infection and the role of contamination of the environment in a cystic fibrosis clinic. *The Journal of Hospital Infection*. 1983; **4**: 31-40.
180. Festini F, Taccetti G, Mannini C, Campana S, Mergni G, Vignoli N, Allegretti N, Ravenni N, Cocchi P, Neri S, Repetto T, de Martino M. Patient risk of contact with respiratory pathogens from inanimate surfaces in a cystic fibrosis outpatient clinic. A prospective study over a four-year period. *Pediatric Pulmonology*. 2007; **42**: 779-84.
181. Wainwright CE, France MW, O'Rourke P, Anuj S, Kidd TJ, Nissen MD, Sloots TP, Coulter C, Ristovski Z, Hargreaves M, Rose BR, Harbour C, Bell SC, Fennelly KP. Cough-generated aerosols of Pseudomonas aeruginosa and other Gram-negative bacteria from patients with cystic fibrosis. *Thorax*. 2009; **64**: 926-31.
182. Moore JE, McIlhatton B, Buchanan J, Gilpin D, Shaw A, Hall V, Murphy PG, Elborn JS. Occurrence of Burkholderia cepacia in the hospital environment. *Ir J Med Sci*. 2002; **171**: 131-3.



183. Ensor E, Humphreys H, Peckham D, Webster C, Knox AJ. Is Burkholderia (Pseudomonas) cepacia disseminated from cystic fibrosis patients during physiotherapy? *The Journal of Hospital Infection*. 1996; **32**: 9-15.
184. Pegues DA, Schidlow DV, Tablan OC, Carson LA, Clark NC, Jarvis WR. Possible nosocomial transmission of Pseudomonas cepacia in patients with cystic fibrosis. *Archives of Pediatrics & Adolescent Medicine*. 1994; **148**: 805-12.
185. Whittington AM, Whitlow G, Hewson D, Thomas C, Brett SJ. Bacterial contamination of stethoscopes on the intensive care unit. *Anaesthesia*. 2009; **64**: 620-4.
186. Hutchinson GR, Parker S, Pryor JA, Duncan-Skingle F, Hoffman PN, Hodson ME, Kaufmann ME, Pitt TL. Home-use nebulizers: a potential primary source of Burkholderia cepacia and other colistin-resistant, gram-negative bacteria in patients with cystic fibrosis. *Journal of Clinical Microbiology*. 1996; **34**: 584-7.
187. Estivariz CF, Bhatti LI, Pati R, Jensen B, Arduino MJ, Jernigan D, Lipuma JJ, Srinivasan A. An outbreak of Burkholderia cepacia associated with contamination of albuterol and nasal spray. *Chest*. 2006; **130**: 1346-53.
188. Brzezinski LX, Riedi CA, Kussek P, Souza HH, Rosario N. Nebulizers in cystic fibrosis: a source of bacterial contamination in cystic fibrosis patients? *Jornal Brasileiro de Pneumologia*. 2011; **37**: 341-7.
189. Festini F, Taccetti G, Galici V, Neri S, Bisogni S, Ciofi D, Braggion C. A 1-m distance is not safe for children with cystic fibrosis at risk for cross-infection with Pseudomonas aeruginosa. *Am J Infect Control*. 2010; **38**: 244-5.
190. Bell SC, Robinson PJ, Fitzgerald DA. 2008. Cystic Fibrosis Standards of Care Australia. http://www.thoracic.org.au/imagesDB/wysiwyg/CF_standardsofcare_Australia_2008.pdf. Accessed: July 14 2013.
191. Standards of care for cystic fibrosis in New Zealand. 2010 http://www.thoracic.org.au/imagesDB/wysiwyg/CF_StandardsofCare_NZ2010.pdf. Accessed: July 14 2013





The Thoracic Society of Australia and New Zealand Ltd

Suite 405, Level 4,
5 Hunter Street,
Sydney NSW 2000

P +61 2 9222 6200

F +61 2 9221 0438

E info@thoracic.org.au

www.thoracic.org.au